

LETTERS TO THE EDITOR

Hypocomplementemic paraneoplastic vasculitis: a rare case of b-cell malignancy

Atakan Sogur O¹, Bulut Gokten D¹, Mercan R¹

Dear Editor,

Pemphigus foliaceus (PF) is an autoimmune skin disorder characterized by superficial vesicle formation and is rarely reported among the cutaneous manifestations associated with non-Hodgkin lymphoma (NHL)¹.

Urticarial vasculitis has been classified into two subgroups based on complement levels: normocomplementemic urticarial vasculitis and hypocomplementemic urticarial vasculitis². Patients with normal complement levels have normocomplementemic urticarial vasculitis, typically with better prognosis, while those with low levels have hypocomplementemic forms, carrying higher complication risk, especially with systemic involvement³. Hypocomplementemia, reflecting chronic antigenic stimulation and autoreactive B-cell activity, signals systemic autoimmunity and risk of lymphomagenesis⁴. In this case, no urticarial lesions were observed, making it a rare atypical presentation of hypocomplementemic urticarial vasculitis syndrome (HUVS) and highlighting the need to consider underlying malignancy, as a B-cell lymphoproliferative disorder was identified.

A 24-year history pemphigus patient presented with recurrent heat, discoloration, and redness of the right hand, reporting two prior similar episodes. On systemic examination, blood pressure was 122/70 mmHg, pulse 80 beats/min, respiratory rate 20/min, and body temperature 36.4°C. Physical examination revealed an erythematous-violaceous lesion with well-defined but irregular borders over the thenar region of the right hand. No urticarial lesions were observed. Hepatomegaly was noted, but there was no splenomegaly. Laboratory tests demonstrated decreased complement levels (C3: 0.73 g/L, C4: 0.01 g/L) and thrombocytopenia (platelet count: 32×10³/μL). Antinuclear antibody (ANA) testing was positive at titers of 1/100–1/320 (1+) with a fine speckled cytoplasmic pattern. Extractable nuclear antigen (ENA) panel revealed no pathological

findings. Hemoglobin level was 14.4 g/dL, and white blood cell (WBC) count was 20.96×10³/μL. Renal and liver function tests were within normal limits. Based on these findings, hypocomplementemic vasculitis was considered, and the patient was referred to the hematology department for further evaluation due to the possibility of a paraneoplastic condition. Subsequent investigations revealed a CD5-negative, CD19-positive B-cell lymphoproliferative disorder. Genetic testing for hereditary cancer-associated mutations did not identify any pathogenic or likely pathogenic variants.

HUVS, also known as McDuffie syndrome, is a rare disorder first described by McDuffie and colleagues in 1973⁵. It is classified as primary (idiopathic, usually not systemic) or secondary (associated with systemic inflammation and often chronic)⁶. The secondary form is characterized by low serum complement, autoantibodies, and immune deposition at the dermoepidermal junction, resembling a lupus band and overlapping with features of systemic lupus erythematosus (SLE)⁷. HUVS has been linked to malignancies, especially Hodgkin and non-Hodgkin lymphomas, though it remains unclear whether it is a cause or consequence of the malignancy⁸. A case report in the literature described a patient with rare hypocomplementemic urticarial vasculitis syndrome presenting as a paraneoplastic manifestation following a diagnosis of chronic lymphocytic leukemia⁹. An erythematous-violaceous lesion with irregular but well-defined borders was noted on the right thenar region, showing central color accentuation with a slightly edematous, rough surface. The appearance of the lesion was consistent with vasculitis (see figure 1). At presentation and previously, the patient had no urticarial lesions, distinguishing this case from others. Notably, marked C3 and C4 hypocomplementemia was observed during the acute episode.

Thrombocytopenia can be an indicator of malignancy. In patients with thrombocytopenia, hematologic malignancies should be considered and ruled out¹⁰. In the present case, laboratory tests during the acute episode revealed thrombocytopenia in addition to complement deficiency.

This case report highlights the potential association between a rare HUVS case presenting with atypical clinical features and an accompanying lymphoproliferative disorder. In the presence of underlying autoimmunity,

¹ Department of Rheumatology, Namik Kemal University Faculty of Medicine, Tekirdag, Türkiye

Submitted: 01/09/2025

Accepted: 03/09/2025

Correspondence to: Omer Atakan Sogur
E-mail: atakan.371@hotmail.com



Figure 1. Erythematous-violaceous lesion over the thenar region of the right hand. The lesion presents with well-defined but irregular borders, a slightly edematous and roughened surface and more intense coloration at the center, consistent with vasculitic changes.

hematologic malignancies should be carefully investigated and ruled out when hematologic abnormalities such as hypocomplementemia and thrombocytopenia accompany cutaneous vasculitic lesions.

REFERENCES

1. Prodanovic EM, Korman NJ. A case of non-Hodgkin's lymphoma-associated pemphigus foliaceus and use of rituximab for treatment. *J Dermatolog Treat.* 2008;19:368-370. <https://doi.org/10.1080/09546630802032603>
2. Jachiet M, Le Quellec A, Deroux A, Godmer P, Ebbo M, Astudillo L, et al. Clinical-biological spectrum and therapeutic management of hypocomplementemic urticarial vasculitis: data from a French nationwide study on 57 patients. *Arthritis Rheumatol.* 2014;66:S1212-S3.
3. Buck A, Christensen J, McCarty M. Hypocomplementemic urticarial vasculitis syndrome: a case report and literature review. *J Clin Aesthet Dermatol.* 2012;5:36.
4. Nocturne G, Pontarini E, Bombardieri M, Mariette X. Lymphomas complicating primary Sjögren's syndrome: from autoimmunity to lymphoma. *Rheumatology (Oxford).* 2021;60:3513-3521. <https://doi.org/10.1093/rheumatology/kez052>
5. McDuffie F, Sams W Jr, Maldonado J, Andreini P, Conn D, Samayoa E, editors. Hypocomplementemia with cutaneous vasculitis and arthritis: possible immune complex syndrome. *Mayo Clin Proc.* 1973.
6. Filosto M, Cavallaro T, Pasolini G, Broglio L, Tentorio M, Cottelli M, et al. Idiopathic hypocomplementemic urticarial vasculitis-linked neuropathy. *J Neurol Sci.* 2009;284:179-181. <https://doi.org/10.1016/j.jns.2009.03.027>
7. Firestein GS, Budd RC, Harris ED Jr, et al., editors. *Kelly's textbook of rheumatology.* 8th ed. Philadelphia: Saunders; 2008. Vol II. Part 13, Chapter 85: Urticarial vasculitis.
8. James WD, Elston D, Berger T. *Andrew's diseases of the skin e-book: clinical dermatology.* Elsevier Health Sciences; 2011.
9. Pereira HP, Alves MP, Coutinho IA, Carrapatoso I, Todo-Bom A. Hypocomplementemic urticarial vasculitis syndrome: a look beyond urticarial lesions. *Adv Dermatol Allergol.* 2023;40:571-573. <https://doi.org/10.5114/ada.2023.130474>
10. Jinna S, Khandhar PB. Thrombocytopenia. 2019.